

Are Mammograms failing us as a diagnostic tool? Many women are refusing to get their annual mammogram. The thought of smashing delicate breast tissue and exposing themselves to carcinogenic radiation in order to see if their breasts are healthy, is causing many women to skip their annual mammogram. This is especially true when considering the error rate of a 40% false negative in postmenopausal women, a 60% false negative rate in women with dense breast tissue and a 80% false positive rate.(1) Out of that 80% false positive, many result in unnecessary, invasive surgical procedures without being informed of far less damaging options. A biopsy of a pre-cancerous lump, breaches the encapsulation, can spread it outside the duct, expose it to a blood supply which can turn it into invasive cancer. This is unnecessary when research has proven these pre-cancerous lumps or Ductal

Carcinoma in Situ, in most cases, can be reversed with an immune supportive protocol(3). When considering that post-mortem statistics show that 40% of the adult female population have DCIS or pre-cancerous cells in their breasts and one out of every eight women will be diagnosed with cancer in their life time these invasive procedures are actually increasing the rate of breast cancer.

Between 1973 and 1981 an in-depth study conducted by the American Cancer Society and National Cancer Institute called the *Breast Cancer Detections and Demonstration Projects* or *BCDDP*, it concluded that: both mammograms and thermogram techniques were lacking in their own way but they chose mammography simply because it could localize a lesion(4). Thermography, however, sees the development of neo-angiogenesis. a bloods supply that feeds a cancerous tumor at a very early stage. Research shows Targeted Ultrasound which is painless and emits no carcinogenetic radiation, has a 99.9% accuracy rate in distinguishing if a lump is fluid filled (as seen with infections i.e. acute cysts) or solid filled with shadow margins (as seen with malignancy), furthermore this study says: "The added value of adjunct with mammography is low"(5), meaning is not necessary.

In 1996, the Head of Radiology at University of California at Berkeley, John Goffman MD., PhD., after 5 years of research, published his findings on how many RADS of radiation it takes to cause [breast] cancer. He felt this information so important, that he published the findings on the cover of his book: *Preventing Breast Cancer*, "Our estimate is that about three-quarters of the current annual incidence of breast-cancer in the United States is being caused by earlier ionizing radiation primarily from medical sources".(6) Goffman's meticulous research suggests that Mammography is actually causing the disease it is suppose to detect.

We must ask ourselves why we are radiating the population for screening just to find a lump when Thermography is safe, non-assaultive and non-invasive more precise diagnostic tool, especially when coupled with Targeted Ultrasound if indicated?

Note: Only with a sophisticated Medical Grade Infrared Camera and Diagnostic Program can accurate reports be generated by a Certified Medical Thermologist. Thus offering Thermography, as a proven, accurate and much less invasive breast screening and monitoring regimen.

Article for Scholarly Review: <u>Time for Women to have a Healthier Choice in Breast Screening</u> How Accurate is Thermography as a Breast Screening Tool?



Thermography is not just a way to find disease, but it is a tool to monitor breast health, without breast compression or radiation. Using the strict procedures outlined in the Marseilles Classification Study, the accuracy rate for both false positive and false negative for *Standardized Medical Thermography*© is 9%.(2) This came from a study in 1976, at the Pasteur Institute, in Marseilles France. In this study, top medical doctors in their respective fields, followed 3,000 women for 5 years using the results of the thermography readings, in addition to histology reports to classify the different ratings of: TH-1 through TH-5 with related risk factors for malignancy. It should be noted that only a biopsy with a histology report can give a 100% positive diagnosis of malignant cells. Unlike Mammography, research has shown that with an Comprehensive Immune

Support Protocol© using Thermography as a monitoring tool 80% of women with Atypical ratings reverse within 3 to 6 months. (7) This has saved hundreds of women from unnecessary invasive procedures.

How to Choose a Thermography Clinic that Follows These Procedures

Here are the factors to establish if a clinic is following *Comprehensive Breast Screening*© with *Standardized Medical Thermography*© criteria:

- 1. Does the clinic take a comprehensive reproductive history and addition to the details of any critical health issue?
- 2. Is the technician qualified to conduct a manual breast exam and recognize palpation of a suspicious lump, dimpling or inverted nipple, Padgett's Disease (breast cancer of the nipple) or inflammatory breast cancer (that doesn't have a lump)? The technician should note the details of their findings on patient's intake forms so changes can be monitored.
- 3. Does the clinic take images before and after performing the *autonomic challenge* as outlined in the Marseilles Procedures? This is important to distinguish neo-angeogenic blood vessels (that set up to feed a tumor), from normal blood vessels. The *autonomic challenge* is acquired by placing the patient's hands on ice for 1 minute. A full set of images, frontal and side views at 45 and 90 degrees on both sides, are taken before and after the *autonomic challenge*. Then the results of the comparison are then noted in the report.
- 4. Does the clinic provide the patient with a comprehensive report that notes the difference in specific features on the thermal images, that need to be monitored when compared to those of the 'normal' breast? This is essential for monitoring improvements in these features. [See the images below for an example of an abnormal reading that has improved with the necessary details noted on the images for comparison.]
- 5. Does the clinic include a comprehensive consultation with each report, to explain the content to the patient? It is important to go over the report in detail to answer the patient's questions, address concerns about specific features, symptoms of complaint or the patient's health in general including stress factors, thereby being able to provide a *Comprehensive Individualize Immune Support Protocol*© to improve the patient's health. This can be

with nutritional supplements to build the immune system, address hormone imbalances and also lifestyle changes to aid in a better sleep or coping with stress, just to mention a few.

- 6. With follow-up thermography sessions and reports, do the findings provide a visual comparison with the previous report so the patient can see if the specific feature of concern has improved or not?
- 7. Last but certainly not least if the patient needs investigative and/or surgical procedures, the professional staff should offer support so they feel comfortable and safe and can make informed decisions, not fear based. As a professional they can only give accurate statistical information and never giving medical advice as a physician.

If the clinic does not follow these procedures the accuracy and credibility of Thermography as a Breast Screening Technique is severely compromised.

Investigative and Surgical Procedures

Early diagnosis before a malignancy has spread to the lymphatic system or other organs has a very good prognosis. However a biopsy of a pre-cancerous condition like DCIS can in fact cause it to advance into a invasive malignancy. Although the Immune Support Protocol has proven to reverse Atypical Breast Thermography or DCIS in 80% of the cases, the progress can be monitored with Thermography (see example below), 20% of cases that do not improve. Some of those are Personal Variants, which means they simply have an asymmetric pattern that has not changed in years and is not progressing or getting worse. For the others in which the features are progressively getting more complex, more prominent, spreading to other quadrants or having other features such as a growing lump, dimpling of the breast or nipple, then other objective means of evaluation is recommended for these patients. Targeted Ultrasound can isolate a lump and tell if it is fluid filled indicating an infection or solid filled. Solid filled could be scar tissue or malignancy the difference being with scar tissue the margins would be clean compared to surrounding tissue in opposed to malignancy where the margins are shadowed indicating it merging into surrounding tissue. If found early a simple lumpectomy is usually sufficient. A sentinel node biopsy is preformed during the surgery before the lump is removed by placing dye into the tumor, it leaks into the connected lymph nodes which are then removed and a histology report of the lump and sentinel nodes are preformed. If there is no lymph involvement chemotherapy is contraindicated. The doctors may suggest or even insist on radiation due to the surgical procedure or biopsy having liberated malignant cells. The patient has the right to be informed of the damage to lungs and heart (if the left breast is involved) through the use of radiation treatment. These stray malignant cells, if in fact there are any can be addressed by using Turkey Tail Mushrooms to build the immune system. The patient will also be tested for Estrogen, Progesterone receptor negative or positive. The Oncologist may recommend some form of Tamoxifin which is normally prescribed for an Estrogen and/or Progesterone positive condition. The patient also has the right to know that this can cause uterine cancer. Other options for lowering the estrogen or progesterone is with Calcium'd Glucarate, which increases Phase 2 liver function to clean out excessive hormones and has been use to replace these drugs with no side effects. Before during and after surgical procedures the use of the homeopathic, Arnica Montana for trauma and injury is very helpful in the healing process.

Example of Thermal Images

Below are images of a patient with Ductal Carcinoma in Situ (DCIS) before and after following an *Comprehensive Individualize Immune Support Protocol*©(7). Pictured below are the proper notations of temperatures on the BEFORE image, comparing the atypical feature (points 1-3) in the right breast with the left or 'normal' features in the left breast. In the AFTER image the atypical feature is gone (see images below).




Image of a patient with a bilateral mammary duct infection, seen as diffuse hyperthemic patterns in the periareolar (nipple) region bilaterally. This atypical but benign condition can be address with the Healing a Mammary Duct Infection Protocol (see image above).






Image of a patient with 30 year-old augmentation implants that are decomposing and leaking into the lymphatic nodes and inflaming the liver (see image above). Thermography can help to reveal other reasons a patient is not feeling well. Not all health issues can be alleviated with nutritional protocols, but may need surgical procedures in addition, as in this case.


Thermography Images Shows Benign, Atypical and Abnormal Ratings of Breasts



Benign Thermography - TH-2 with symmetric vascular-like features.



Atypical Thermography - TH-3 with 1 asymmetric vascular-like feature.



Abnormal Thermography - TH-5 of the right breast with a complex vascuar-like feature (points 1&2 in the right view) indicating neo-angiogenesis, a blood supply associated with malignancy. A TH-5 rating has 3 criteria: 1. A complex vascular feature, 2. This feature is seen in 2 quadrants, cranial lateral and caudal lateral and 3. It is 1.4°C above of base emission levels which is considered an anarch sign. The left breast is rated: TH-2 or benign with no thermal features of concern, the hotest point noted for comparison.

Conclusion

Comprehensive Breast Screening® with Standardized Medical Thermography®, using Target Ultrasound for follow-up investigation when needed, is a far more accurate and less harmful technique for breast screening and monitoring women's breast health, and needs to be made available to all women.

It is time to stand up for our right to choose a more accurate and less harmful tool to screen and improve the health of our breasts.

Acknowledgements

The author would like acknowledge two prominent Bay Area Surgeons: Dr. Ann Verecoutere and Dr. William Small. Thank you for the kindness and respect you have shown my patients. Somehow thank you is not enough for the patience and willingness to answer numerous questions in order to expand my knowledge. This service has been instrumental in giving an indepth understanding of the scope of investigative tools, surgical

procedures, and breast oncology in general. Also invaluable to me is my friendship with and the professional support of Dr. Lauren Eskanazi, author of *Reconstructing Aphroditie* and a phenominal Plastic Surgeon as well as artist in breast reconstruction.

About the Author



Article by Nancy Gardner-Heaven, Technician of Medical Thermology, Certified at Auburn University by the American Academy of Thermology. Nancy is a pioneer in the field of Medical Thermology offering *Comprehensive Breast Screenings*© using *Standardized Medical Thermography* © for 25 years. Research from her immune boosting protocols has proven to reverse DCIS and other Atypical Features in 80% of her patients. This study was presented at Auburn University at the American Academy of Thermology's Medical Conference(3) in 2003 and published in the Townsend Letter in 2004. Her published research has been instrumental in saving thousands of women from unnecessary surgical procedures.

Nancy continues to improve this protocol as more substantial research surfaces. She is the founder of *Comprehensive Breast Screening*© with *Standardized Medical Thermography*© and has been using this technique since 1999. Comprehensive Breast and Full Body Screening is available in the greater San Francisco Bay Area as well as the Big Island and Kauai. See website: www.HeatlhyBreasts.Info

- (1) American Cancer Society Annual Statistics of Mammography Accuracy Rate.
- (2) Marseilles Classification Study at the Pasteur Institute in Marseilles, France 1976.
- (3) Abstract on Research of Immune Support Protocols Reversing Atypical Features presented at Auburn University, Published in the Townsend Letter.
- (4) Article: Infrared Imaging: Making Progress in Fulfilling Its Medical Promise Published in IEEE Engineering in Medicine and Biology Magazine by Drs Robert Elliott and Joanathan F Head Volume 21, Number 6 Nov/Dec 2002.
- (5) Article: Accuracy and Value of Breast Ultrasound for Primary Imaging Evaluation of Symptomatic Women 30-39 Years of Age Published in: American Journal of Roentgenology Diagnostic Imaging Related Sciences by November 2012, Volume 199, #5
- (6) Preventing Breast Cancer The Story of a Major, Proven, Preventable Cause of This Disease by John W. Goffman, MD, Ph.D. Second Edition 1996 C.N.R. Book Division Published by: Committee for Nuclear Responsibility, Inc.
- (7) Abstract Atypical Breast Thermology Case Study of Protocol Reversing DCIS and Other Atypical Breast in 80% of Patients, published in the Townsend Letter for Doctors and Patients- November 2004

Contact Information:
Nancy Gardner-Heaven
P.O. Box 891
Knights Ferry, CA 95361-2091

Phone (209) 881-3044

Intellectual Property by Nancy Gardner-Heaven All Rights Reserved Internationally @ 2-2-19 No portion of this article may be published without written consent and acknowledgement of the author.